

Announcement:

- Office hour: Tuesday 1:30 - 2:30 PM
(Zoom)


Today:

- Bisection Method
- Intro to fixed point problems

HW questions?

Bisection Method

Picture:



EEx/ $f(x) = x^2 - 2$
4 iterations of B.M. on $[1, 2]$

Solu Step 0 $f(x)$ is continuous (polynomial)

$$\cdot f(1) = -1 < 0$$

$$\cdot f(2) = 2 > 0 \quad \checkmark$$

k	a_k	b_k	p_k	$f(a_k)$	$f(b_k)$	$f(p_k)$
1	1	2	$\frac{3}{2}$	-1	2	$\frac{1}{4}$
2	1	$\frac{3}{2}$	$\frac{5}{4}$	-1	$\frac{1}{4}$	$-\frac{7}{16}$
3	$\frac{5}{4}$	$\frac{3}{2}$	$\frac{11}{8}$	$-\frac{7}{16}$	$\frac{1}{4}$	$-\frac{23}{64}$
4	$\frac{11}{8}$	$\frac{3}{2}$	$\frac{23}{16}$			

Error bound

$$|p_n - p| \leq \frac{b-a}{2^n}, \quad n \geq 1$$

EEx/ $x^3 + x - 4 = 0$

B.M. on $[1, 4]$

of iterations required for an accuracy
of 10^{-3}

Soln Step 0 (note): $f(x) = x^3 + x - 4$
is cont. (polynomial)

$$f(1) < 0$$

$$f(4) > 0$$

$$|p_n - p| \leq \frac{b-a}{2^n} < 10^{-3}$$

$$\frac{3}{2^n} < 10^{-3}$$

$$n > 11.6$$

\Rightarrow Need no more than 12 iterations

Fixed point formulations

Ex $f(x) = x^3 - 2x + 1$

For each $g(x)$, show that

$$f(p) = 0 \text{ if and only if } g(p) = p$$

a) $g(x) = \frac{1}{2}(x^3 + 1)$

b) $g(x) = \frac{2}{x} - \frac{1}{x^2}$

c) $g(x) = \sqrt{2 - \frac{1}{x}}$

d) $g(x) = -\left(\sqrt[3]{1 - 2x}\right)$

Solv

a) $f(p) = 0 \Leftrightarrow p^3 - 2p + 1 = 0$

$$\Leftrightarrow 2p = p^3 + 1$$

$$\Leftrightarrow p \in \frac{1}{2}(p^3 + 1)$$

$$g(p) = p$$

b) $f(p) = 0 \Leftrightarrow p^3 - 2p + 1 = 0$

$$\Leftrightarrow p^3 = 2p - 1$$

$$p = \frac{2}{p} - \frac{1}{p^2} \quad (\text{OK since } f(p) = 0, p \neq 0)$$

$$\Leftrightarrow g(p) = p$$

c), d) similar